



**NON-HOSPITAL MEDICAL AND SURGICAL FACILITIES
ACCREDITATION PROGRAM**

College of Physicians and Surgeons of British Columbia

Obesity

GUIDELINE

You may download, print or make a copy of this material for your non-commercial personal use. Any other reproduction in the whole or part requires written permission from the College of Physicians and Surgeons of British Columbia and the material must be credited to the College.

Approved: May 2006

Revised: May 2017, December 30, 2017 (Program name change only)

Originating committee: Non-Hospital Medical and Surgical Facilities Program Committee

Preamble

The Non-Hospital Medical and Surgical Facilities Accreditation Program (NHMSFAP) Committee develops professional guidelines to assist physicians in meeting high standards of medical practice and conduct. A guideline reflects a recommended course of action established based on the values, principles and duties of the medical profession. Guidelines attempt to define practices that meet the needs of patients in most circumstances and are not a replacement for clinical judgment. As a result, situations may arise in which deviations from these guidelines may be appropriate and physicians may exercise reasonable discretion in their decision-making based on the guidance provided.

The NHMSFAP Committee recognizes that obesity is a major contributor to chronic diseases; however, there is a need for access to high-quality surgical care. The decision to recommend surgery for obese patients in the non-hospital setting requires comprehensive evaluation of the patient's ASA classification, medical history, comorbidities and lifestyle habits to assess weight-related health issues and the impact these will have on a proposed surgery and anesthetic.

Patients with ASA 1 and 2 can be considered for surgery in a non-hospital facility. Patients with ASA 3 may be accepted under certain circumstances.

Practice Guidelines

Obese patients must be carefully screened prior to a proposed surgery and anesthesia.

Patients are screened using: Body mass index (BMI) = weight in kg/height in metres²

1. BMI \geq 30 and $<$ 40 – Obesity Classification I (30–34.9) and II (35–39.9)

Consideration of surgery and anesthesia for patients with a BMI \geq 30 and $<$ 40 the following conditions should be met:

- a. Comorbid conditions are well-controlled and unlikely to add significant risk to the procedure or anesthetic; and
- b. The proposed surgery and anesthesia are unlikely to aggravate or precipitate significant changes in these conditions.

2. BMI \geq 40 – Obesity Classification III (Morbid Obesity)

The decision to perform surgery on a patient with morbid obesity of BMI \geq 40 should take into account the health risks and comorbidities associated with the planned surgery and the impact those conditions will have on the patient outcome.

Consideration of surgery and anesthesia for patients with a BMI \geq 40 the following conditions should be met:

- a. Patient presents with no other severe systemic disease with substantive functional limitation other than morbid obesity of BMI \geq 40;
- b. Patient's morbid obesity is unlikely to add significant risk to the procedure or anesthetic;
- c. The patient must have had a recent recorded history, physical examination, and appropriate laboratory and diagnostic investigations within 60 days of the surgery;

- d. The proposed anesthesia should be limited to topical, local or peripheral nerve block anesthesia, or anesthesia with single-agent oral sedation if needed; however,
- e. Under certain circumstances, at the discretion of the preoperative and intra-operative anesthesiologists, the delivery of a higher level of anesthesia (i.e. IV sedation/analgesia, regional block or general anesthetic) may be acceptable; the plan must be consistent to that set out in the preoperative anesthetic consult;
- f. If considering a higher level of anesthesia (i.e. IV sedation/analgesia, regional block or general anesthetic), a documented preoperative anesthetic consultation must be completed before admission for the surgical procedure, not more than 28 days before surgery and at minimum, one day prior to surgery. The consult must include the possible impact of the surgery, plan for anesthesia and the post-operative course;
- g. All patients with BMI ≥ 40 who require **unplanned** IV sedation/analgesia, regional block or general anesthetic during the course of any surgery must be reported by the medical director to the College using the NHMSFAP reportable incident form; and
- h. In all circumstances, **planned or unplanned**, for which a higher level of anesthesia is required an anesthesiologist must be dedicated to the patient at all times throughout the conduct of all IV sedation/analgesia, regional block or general anesthetic until the patient is transferred to the post anesthesia recovery.

3. BMI ≥ 45 – Obesity Classification III (Morbid Obesity)

Patients with a BMI ≥ 45 are **not suitable for surgery under any circumstance** in the non-hospital setting.

4. General recommended principles of care for the obese patient include but are not limited to:

- a. Specialized equipment for treating obese patients (e.g. special seating/stretchers/OR table/transfer mats/toilet/sinks and monitoring equipment)
- b. Planned management of potential surgical complications
- c. Follow-up care and outcome reporting

References

American Society of Anesthesiologists Clinical Information [Internet]. Schaumburg, IL: American Society of Anesthesiologists; 2017. ASA physical status classification system; [approved 2014 Oct 15; cited 2017 Feb 6]; 1 screen. Available from: <https://www.asahq.org/resources/clinical-information/asa-physical-status-classification-system>

Dobson G, Chong M, Chow L, Flexman A, Kurrek M, Laflamme C, Lagace A, Stacey S, Thiessen B. Guidelines to the practice of anesthesia - revised edition 2017. Can J Anaesth [Internet]. 2017 Jan [cited 2017 Feb 6];64(1):65-91. Available from: http://www.cas.ca/English/Page/Files/97_Guidelines-2016.pdf

NICE: National Institute for Health and Care Excellence (UK). Obesity: identification, assessment and management [Internet]. London: NICE; 2014. [cited 2017 Feb 6]. 64 p. (Clinical guideline: CG 189). Available from: <https://www.nice.org.uk/guidance/cg189/chapter/1-recommendations>

Sharma A, Kushner R. EOSS: Edmonton obesity staging system [Internet]. Edmonton: Alberta Health Services; University of Alberta; 2009. [cited 2017 Feb 6]. 1 p. Available from:
<http://www.drsharma.ca/wp-content/uploads/edmonton-obesity-staging-system-staging-tool.pdf>

World Health Organization Global Database on Body Mass Index [Internet]. Geneva, Switzerland: WHO; 2006. BMI classification [updated 2017 Feb 6; cited 2017 Feb 6]; 1 screen. Available from:
http://apps.who.int/bmi/index.jsp?introPage=intro_3.html